

B_K from improved staggered quarks *

E.Gámiz^a, S.Collins^a, C.T.H.Davies^a, J.Shigemitsu^b, M.Wingate^c, HPQCD and UKQCD collaborations.

^aDepartment of Physics & Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK.

^bPhysics Department, The Ohio State University, Columbus, OH 43210, USA.

^cInstitute for Nuclear Theory, University of Washington, Seattle, WA 98115, USA.

We compare calculations of B_K with improved staggered quarks (HYP, Asqtad) and demonstrate the improved scaling behaviour that this gives rise to over previous calculations with unimproved staggered quarks. This enables us to perform the calculation of B_K on the MILC dynamical configurations ($n_f = 2 + 1$), for which we give preliminary results.

1. Introduction

Indirect CP violation in the system of neutral kaons is parametrized by ε_K . Experimentally it is a very well measured quantity. On the other hand, its theoretical expression depends on certain CKM matrix elements about which we would like to obtain information and on the hadronic matrix element between K^0 and $\bar{K}^0$ of the effective hamiltonian describing processes with $\Delta S = 2$,

$$H_{eff}^{\Delta S=2} = C_{\Delta S=2}(\mu) \int d^4x Q_{\Delta S=2}(x) \quad (1)$$

with

$$Q_{\Delta S=2}(x) = [\bar{s}_\alpha \gamma_\mu d_\alpha]_{V-A}(x) [\bar{s}_\beta \gamma^\mu d_\beta]_{V-A}(x). \quad (2)$$

The matrix element of this operator is usually normalized by its VIA value, defining B_K as the ratio

$$B_K(\mu) \equiv \frac{\langle \bar{K}^0 | Q_{\Delta S=2}(\mu) | K^0 \rangle}{\frac{8}{3} \langle \bar{K}^0 | \bar{s} \gamma_\mu \gamma_5 d | 0 \rangle \langle 0 | \bar{s} \gamma_\mu \gamma_5 d | K^0 \rangle}. \quad (3)$$

The theoretical error associated with the calculation of B_K is the main source of uncertainty when one tries to use the experimental value of ε_K to constraint the CKM matrix, so improvement in the determination of this parameter is crucial

to gain information about the unitarity triangle. The lattice calculations have an important role to play in such reduction of errors.

2. Calculation of B_K using staggered fermions

The staggered quark formulation has the advantage of conserving some chiral symmetry at nonzero lattice spacing. This residual symmetry forbids the mixing with operators of different chirality leading to a much simpler renormalization process. In addition, dynamical calculations with staggered quarks are feasible with present computers.

The value of B_K used by phenomenologists at present in CKM studies was obtained by the JLQCD collaboration [1] using unimproved staggered quarks in the quenched approximation. The value given in [1] is $B_K^{NDR}(2 \text{ GeV}) = 0.628(42)$. In spite of being considered the benchmark of the calculations of B_K , the determination by the JLQCD collaboration has several drawbacks. First, it presents large scaling corrections, so the extrapolation to the continuum is not as reliable as it could be with a better scaling behaviour. The main uncertainty is quenching, whose possible impact was estimated in [2] using chiral perturbation theory to be of the order of 15%. The $SU(3)$ breaking effects are not incorporated in the calculation either, but this prob-

*Talk presented by E. Gámiz at *Lattice 2004*, Fermilab, 21–26 June 2004.




Figure 1. Scaling of $B_K^{NDR}(2\text{GeV})$ with a for improved staggered actions compared to the JLQCD unimproved staggered results. The stars are values calculated by renormalizing the unimproved bare matrix with the HYP Wilson coefficients. See text for a discussion of renormalization.

ably has a minor effect [2]. We show here that the large scaling violation can be corrected by using improved staggered fermions. On the other hand, the dominant error, associated to quenching, can be reduced by performing dynamical simulations. And, finally, the use of kaons made of two non-degenerate quarks can account for the $SU(3)$ breaking effects. This three-fold improvement of the staggered determination is the goal of our calculation.

3. Improved staggered actions

We first check the impact of using improved actions in the quenched approximation. We study two such actions, the HYP action [3] and the Asqtad action [4]. Fig. 1 shows our results for unimproved staggered, Asqtad and HYP actions along with the JLQCD results, for several values of the lattice spacing.

Two comments are in order in relation with Fig. 1. First of all, we match kaon masses at a given β to those of the JLQCD collaboration to make a clear comparison. Secondly, the Wil-




Figure 2. B_K^{bare} as a function of the timeslice ($m_{sea} = 0.01/0.05$).

son coefficients needed in the calculation of the renormalized B_K with the Asqtad action are not available at present [5]. Hence, as a first estimate, we use the Wilson coefficients for HYP [6] in the Asqtad case in Fig. 1. This is incorrect, but large differences are unlikely because the 1-gluon vertex is the same for HYP and the $\overline{Fat7}$ smearing which goes into Asqtad.

Fig.1 shows that discretization errors are much smaller with the improved actions. This improvement is largely due to the change in renormalization factors, not to a change in the bare matrix elements. The contribution of several matrix elements confuses the picture but one way to show this is given in Fig. 1. There, the stars correspond to the values of $B_K^{NDR}(2\text{GeV})$ obtained with the unimproved bare matrix elements and the HYP renormalization coefficients.

We expect improved scaling to survive unquenching and therefore allow us to perform reliable dynamical calculations with only a few values of the lattice spacing and even obtain useful information from a single point simulation.

4. Dynamical effects in the calculation of B_K

The next step is to incorporate dynamical effects. We performed a dynamical calculation of B_K within the Asqtad action, using the configurations from the MILC collaboration with $n_f = 2 + 1$ dynamical flavours [7]. Two ensembles at $a = 0.125 fm$ are used: $m_{sea} = 0.01/0.05$ and $m_{sea} = 0.02/0.05$. From the hadron masses determined after the simulation, the strange quark mass, am_s , was found to be 0.04. We make kaons on the configurations from degenerate quarks of bare mass $m_s/2 = 0.02$.

Fig. 2 shows an excellent plateau when B_K^{bare} is plotted as a function of the lattice timeslice.

In the absence of the correct renormalization coefficients, we again estimate $B_K^{NDR}(2GeV)$ from the HYP Wilson coefficients for unimproved glue. The results we obtained in this way for the two different values of the light quark mass, including only statistical errors, together with the result quoted by the JLQCD collaboration and its corresponding uncertainty, are depicted in Fig. 3.

From the results in Fig. 3 it seems likely that B_K from full QCD will be close to the quenched result. There is also some sign that $B_K^{NDR}(2GeV)$ is falling with the dynamical quark mass.

5. Summary

We found that the large scaling corrections that affected previous staggered calculations of B_K are reduced by using improved actions (HYP, Asqtad) in the quenched approximation. This improved behaviour allows us to perform a reliable dynamical calculation of this non-perturbative parameter on the MILC configurations. We give a preliminary estimate for $B_K^{NDR}(2GeV)$ in Fig. 3.

In order to get a final dynamical value for $B_K^{NDR}(2GeV)$ and to clear up the role of dynamical effects in this quantity, we must redo our calculation using the correct renormalization coefficients, obtain results for another lattice spacing and eventually, although its effect is less important, consider physical kaons instead of kaons made up of two degenerate quarks with $m_s/2$.




Figure 3. An estimate of $B_K^{NDR}(2GeV)$ from dynamical lattice QCD. The x axis corresponds to the light sea quark mass divided by the (real) strange quark mass. The lines represent the quenched result from [1].

Acknowledgements: E.G. is indebted to the European Union for a Marie Curie Intra-European Fellowship. The work is also supported by PPARC and DoE.

REFERENCES

1. S. Aoki *et al.* [JLQCD Collaboration], Phys. Rev. Lett. **80** (1998) 5271 [hep-lat/9710073].
2. S. R. Sharpe, Nucl. Phys. Proc. Suppl. **53** (1997) 181 [hep-lat/9609029].
3. A. Hasenfratz and F. Knechtli, Phys. Rev. D **64** (2001) 034504 [hep-lat/0103029].
4. S. Naik, Nucl. Phys. B **316** (1989) 238; G. P. Lepage, Phys. Rev. D **59** (1999) 074502 [hep-lat/9809157]; K. Orginos, D. Toussaint and R. L. Sugar [MILC Collaboration], Phys. Rev. D **60** (1999) 054503 [hep-lat/9903032].
5. T. Becher and K. Melnikov, in progress.
6. W. J. Lee and S. Sharpe, Phys. Rev. D **68** (2003) 054510 [hep-lat/0306016].
7. C. W. Bernard *et al.*, Phys. Rev. D **64** (2001) 054506 [hep-lat/0104002].